



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Solution by G. B. M. ZERR, A. M., Ph. D., The Temple College, Philadelphia, Pa., and J. R. HITT, Coronal Institute, San Marcos, Tex.


Project the given triangle into an equilateral triangle, side= a ; let G be the center of gravity, CH or $CM=x$.

$$\text{Then } BK = \frac{a(a-2x)}{2a-3x}, \quad AK = \frac{a(a-x)}{2a-3x}, \quad AH = a-x.$$

$$\text{Area } AKH = \frac{\Delta(a-x)^2}{a(2a-3x)}, \quad BN = \frac{2ax-a^2}{3x-a},$$

$$CN = \frac{ax}{3x-a}.$$

$$\text{Area } CMN = \frac{\Delta x^2}{a(3x-a)}, \quad \text{area } ABNM = \Delta \left[1 - \frac{x^2}{a(3x-a)} \right].$$



$$\begin{aligned} p &= \left[\int_0^{\frac{1}{2}a} \frac{(AKH)^3 dx}{\Delta^3} + \int_{\frac{1}{2}a}^a \frac{(ABNM)^3 dx}{\Delta^3} \right] / \int_0^{\frac{1}{2}a} dx \\ &= \frac{2}{a} \int_{\frac{1}{2}a}^a \left[1 - \frac{x^2}{a(3x-a)} \right]^3 dx + \frac{2}{a^4} \int_0^{\frac{1}{2}a} \frac{(a-x)^3 dx}{(2a-3x)^3} \\ &= \frac{2}{a} \int_{\frac{1}{2}a}^a \left[1 - \frac{x^2}{a(3x-a)} \right]^3 dx + \frac{2}{a^4} \int_{\frac{1}{2}a}^a \frac{x^6 dx}{(3x-a)^3} \\ &= \frac{2}{a} \int_{\frac{1}{2}a}^a dx - \frac{6}{a^2} \int_{\frac{1}{2}a}^a \frac{x^2 dx}{3x-a} + \frac{6}{a^3} \int_{\frac{1}{2}a}^a \frac{x^4 dx}{(3x-a)^2} \\ &= \frac{1}{2} \left(\frac{2}{3} - \frac{2}{3} \log 2 \right). \end{aligned}$$

116. Proposed by the late ENOCH BEERY SEITZ.

The average area of the quadrilateral formed by joining four random points on the surface of a circle, radius a , is $4a^2/3\pi$.


Solution by G. B. M. ZERR, A. M., Ph. D., Professor of Chemistry and Physics. The Temple College, Philadelphia, Pa.

Let a =radius of given circle, A =its area, Δ =required average, Δ' =the average area when the four points are taken on both the circle A and a concentric annulus B , Δ_1 =the average area when three points are taken on A and one on B .

$$\text{Then } (\Delta' - \Delta)A = 4B(\Delta_1 - \Delta).$$

$$\text{But } \Delta : \Delta' = A : A+B.$$

$$\therefore \Delta' = \frac{(A+B)\Delta}{A}. \quad \therefore \Delta = \frac{4}{5}\Delta_1.$$



Let one point O be on the circumference of the circle, and the other points P , Q , R anywhere on its surface. Let $OP=x$, $OR=y$, OQ

$=z$, $\angle AOD=\theta$, $\angle COD=\phi$, $\angle BOD=\psi$. Then $OA=2a\sin\theta=x'$, $OC=2a\sin\phi=y'$, $OB=2a\sin\psi=z'$. Area $OPQR=\frac{1}{2}xz\sin(\theta-\psi)+\frac{1}{2}yz\sin(\psi-\phi)=u$.

$$\therefore \Delta_1 = \frac{\int_0^\pi \int_0^\theta \int_\phi^\theta \int_0^{x'} \int_0^{y'} \int_0^{z'} uxyz d\theta d\phi d\psi dx dy dz}{\int_0^\pi \int_0^\theta \int_\phi^\theta \int_0^{x'} \int_0^{y'} \int_0^{z'} xyz d\theta d\phi d\psi dx dy dz}$$

$$\begin{aligned} \therefore \Delta &= \frac{4}{5} \Delta_1 = \frac{24}{5\pi^3 a^3} \int_0^\pi \int_0^\theta \int_\phi^\theta \int_0^{x'} \int_0^{y'} \int_0^{z'} uxyz d\theta d\phi d\psi dx dy dz \\ &= \frac{32}{5\pi^3 a^3} \int_0^\pi \int_0^\theta \int_\phi^\theta \int_0^{x'} \int_0^{y'} [x\sin(\theta-\psi) + y\sin(\psi-\phi)] xyz \sin^3 \phi d\theta d\phi d\psi dx dy \\ &= \frac{64}{15\pi^3 a^3} \int_0^\pi \int_0^\theta \int_\phi^\theta \int_0^{x'} [3x\sin(\theta-\psi) + 4a\sin\phi\sin(\psi-\phi)] \sin^2 \phi \sin^3 \phi d\theta d\phi d\psi x dx \\ &= \frac{512a^2}{15\pi^3} \int_0^\pi \int_0^\theta \int_\phi^\theta [\sin\theta\sin(\theta-\psi) + \sin\phi\sin(\psi-\phi)] \sin^2 \theta \sin^2 \phi \sin^3 \phi d\theta d\phi d\psi \\ &= \frac{64a^2}{15\pi^2} \int_0^\pi \int_0^\theta [3(\phi-\theta)(\sin\theta\cos\theta - \sin\phi\cos\phi) + 3(\sin\theta\cos\theta - \sin\phi\cos\phi)^2 \\ &\quad + 2\sin(\theta-\phi)(\sin^3\theta\cos\phi - \cos\theta\sin^3\phi)] \sin^2 \theta \sin^2 \phi d\theta d\phi \\ &= \frac{2a^2}{45\pi^3} \int_0^\pi (105\theta - 72\theta^2 \sin\theta\cos\theta + 84\theta\sin^3\theta - 120\theta\sin^4\theta - 105\sin\theta\cos\theta - 82\sin^3\theta\cos\theta \\ &\quad + 32\sin^5\theta\cos\theta) \sin^2 \theta d\theta = 4a^2/3\pi. \end{aligned}$$

MISCELLANEOUS.

111. Proposed by G. B. M. ZERE, A. M., Ph. D., Professor of Chemistry and Physics in The Temple College, Philadelphia, Pa.

Exhibit $\cos^3\theta\sin^3\theta\sin^2\phi\cos\phi$ as a series of harmonics.

Solution by the PROPOSER.

$$f(\mu, \phi) = \sum_{m=0}^{m=\infty} [A_{0,m}P_m(\mu) + \sum_{n=1}^{n=m} (A_{n,m}\cos n\phi + B_{n,m}\sin n\phi)P_m^n(\mu)].$$

$$\text{Now } f(\mu, \phi) = \mu^3 \sqrt{(1-\mu^2)^3} [\sin^2\phi\cos\phi = \frac{1}{4}\mu^3(1-\mu^2)^{\frac{3}{2}}(\cos\phi - \cos 3\phi)].$$

$$A_{0,m} = \frac{2m+1}{16\pi} \int_{-1}^1 \mu^3(1-\mu^2)^{\frac{3}{2}} P_m(\mu) d\mu \int_0^{2\pi} (\cos\phi - \cos 3\phi) d\phi = 0.$$